Recombinant Human Cochlin/COCH Protein (short isoform, His Tag)

by Elabscience

Catalog Number: PKSH031139

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms COCH-5B2;COCH5B2;DFNA9

Species Human

Expression Host HEK293 Cells
Sequence Glu 25-Gln 550
Accession NP_001128530.1

Calculated Molecular Weight 59.4 kDa

Observed molecular weight 18 kDa

Tag N-His

Properties

Purity > 92 % as determined by reducing SDS-PAGE.
 Endotoxin < 1.0 EU per μg as determined by the LAL method.

Storage Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of


reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4

Reconstitution Please refer to the printed manual for detailed information.

Data



Background

Cochlin, also known as COCH-5B2 and COCH, is a secreted protein which contains one LCCL domain and two VWFA domains. It is an abundant inner ear protein expressed as multiple isoforms. Its function is also unknown, but it is suspected to be an extracellular matrix component. Cochlin and type II collagen are major constituents of the inner ear extracellular matrix, and Cochlin constitutes 70% of non-collagenous protein in the inner ear, the cochlin isoforms can be classified into three subgroups, p63s, p44s and p40s. The expression of cochlin is highly specific to the inner ear. Eleven missense mutation and one in-frame deletion have been reported in the COCH gene, causing hereditary progressive sensorineural hearing loss and vestibular dysfunction, deafness autosomal dominant type 9 (DFNA9). The co-localization of cochlin and type II collagen in the fibrillar substance in the subepithelial area indicate that cochlin may play a role in the structural homeostasis of the vestibule acting in concert with the fibrillar type II collagen bundles. Defects in COCH may contribute to Meniere disease which is an autosomal dominant disorder characterized by hearing loss associated with episodic vertigo.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>