

Recombinant Human GADD45A/DDIT-1 Protein (His & GST Tag)

Catalog No. PKSH031234

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Growth Arrest and DNA Damage-Inducible Protein GADD45 Alpha;DNA Damage-Inducible Transcript 1 Protein;DDIT-1;GADD45A;DDIT1;GADD45
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Arg 165
Accession	P24522
Calculated Molecular Weight	46.2 kDa
Observed molecular weight	46 kDa
Tag	N-His-GST
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

GADD45A is a member of the GADD45 Family, and has been found to associate with several cytoplasmic and nuclear

For Research Use Only

factors and has been implicated in several cellular functions, including MAPK signaling, cell cycle regulation, DNA repair and genomic stability, apoptosis, and immune responses. The GADD45 Family of genes is rapidly induced by different stressors, including differentiation-inducing cytokines, and there is a large body of evidence that their cognate proteins are key players in cellular stress responses. GADD45A protein has been reported to interact with multiple important cellular proteins, including Cdc2 protein kinase, proliferating cell nuclear antigen, p21Waf1/Cip1 protein, core histone protein and MTK/MEKK4, an up-stream activator of the JNK/SAPK pathway, indicating that GADD45A may play important roles in the control of cell cycle checkpoint, DNA repair process, and signaling transduction. GADD45A expression in response to genotoxic stress illustrates a more complex scenario, wherein transcriptional changes operate in concert with mRNA turnover and translational regulation. GADD45A was the first stress-inducible gene determined to be up-regulated by p53 and is also a target for the p53 homologues, p63 and p73. The decreased GADD45A expression is also considered a survival mechanism, as cancer cells without this control can evade the apoptotic pathway leading to increased tumorigenesis. As GADD45A is an essential component of many metabolic pathways that control proliferating cancer cells, it presents itself as an emerging drug target worthy of further investigation.