Recombinant Human S100A12/CAGC Protein

Catalog No. PKSH031241

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	Protein S100-A12;Calcium-binding protein in amniotic fluid 1;Calgranulin-C;Extracellular newly identified RAGE-binding protein;Migration inhibitory factor-related protein 6;S100 calcium-binding protein A12;Calcitermin;S100A12;CGRP;MRP-6;EN-RAGE
Species	Human
Expression Host	E.coli
Sequence	Met 1-Glu 92
Accession	NP_005612.1
Calculated Molecular Weight	10.6 kDa
Observed molecular weight	10 kDa
Tag	None
Bioactivity	Immobilized recombinant human S100A12 at 2 μg/ml (100 μl/well) can bind human AGER with a linear range of 0.032-20 μg/ml.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

S100 proteinis a family of low molecular weight protein found in vertebrates characterized by twoEF-hand calcium-binding motifs. There are at least 21 different S100 proteins, and the name is derived from the fact that the protein is100%soluble in ammonium sulfateat neutralpH. Most S100 proteins are disulfide-linked homodimer, and is normally present in cells derived from theneural crest, chondrocytes, macrophages, dendritic cells, etc. S100 proteins have been implicated in a variety of intracellular and extracellular functions. They are involved in regulation of protein phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, and the inflammatory response. Protein S100-A12, also known as S100 calcium-binding protein A12, Calcium-binding protein in amniotic fluid 1, Calgranulin-C, and S100A12, is a member of the S-101 family. Like the majority of S100 proteins, S100A12 is a dimer, with the interface between the two subunits being composed mostly of hydrophobic residues. The fold of S100A12 is similar to the other known crystal and solution structures of S100 proteins, except for the linker region between the two EF-hand motifs. S100A12 plays an important role in the inflammatory response.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com