A Reliable Research Partner in Life Science and Medicine

Recombinant Human Troponin C/TNNC1 Protein

Catalog No. PKSH031309

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CMD1Z;CMH13;TN-C;TNC;TNNC

Species Human
Expression Host E.coli

SequenceMet 1-Glu161AccessionNP_003271.1Calculated Molecular Weight18.4 kDaObserved molecular weight20 kDaTagNone

Bioactivity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin Please contact us for more information.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile 150 mM NaCl, 10 mM Na2HPO4, pH 7.5


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Troponin I, also known as TNNC1, is part of the troponin complex. This complex contains 3 subunits: troponin I (TnI), troponin T (TnT) and troponin C (TnC). Troponin I is the inhibitory subunit, blocking actin-myosin interactions and

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

thereby mediating striated muscle relaxation. It binds to actin in thin myofilaments to hold the actin-tropomyosin complex in place. Because of it myosin cannot bind actin in relaxed muscle. When calcium binds to the Troponin C it causes conformational changes which lead to dislocation of troponin I and finally tropomyosin leaves the binding site for myosin on actin leading to contraction of muscle.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com