

Recombinant Human Cyclophilin B/PPIB Protein (His Tag)

Catalog No. PKSH031313

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CYP-S1;CYPB;HEL-S-39;OI9;SCYLP
Species	Human
Expression Host	HEK293 Cells
Sequence	Asp 34-Ala 212
Accession	NP_000933.1
Calculated Molecular Weight	22 kDa
Observed molecular weight	22 kDa
Tag	C-His

Properties

Purity	> 96 % as determined by reducing SDS-PAGE.
Storage	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
Reconstitution	Please refer to the printed manual for detailed information.

Data



Background

Thyroxine-binding globulin, also known as T4-binding globulin, Serpin A7 and TBG, is a secreted protein which belongs to the serpin family. TBG is synthesized primarily in the liver as a 54 kDa protein. TBG is genomically a serpin, although it has no inhibitory function like many other members of this class of proteins. TBG binds thyroid hormone in circulation. It is one of three proteins (along with transthyretin and albumin) responsible for carrying the thyroid hormones thyroxine (T4) and 3,5,3'-triiodothyronine (T3) in the bloodstream. Of these three proteins, TBG has the highest affinity for T4 and T3, but is present in the lowest concentration. Despite its low concentration, TBG carries the majority of T4 in serum. Due to the very low serum concentration of T4 and T3, TBG is rarely more than 25% saturated with its ligand.

For Research Use Only

Unlike transthyretin and albumin, TBG has a single binding site for T4/T3. TBG tests are sometimes used in finding the reason for elevated or diminished levels of thyroid hormone.