

Recombinant Human C1QB/C1qB Protein (His Tag)

Catalog No. PKSH031345

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	C1QB
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Ala 253
Accession	NP_000482.3
Calculated Molecular Weight	25 kDa
Observed molecular weight	30 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 94 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, 0.5mM TCEP, 10% glycerol, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 94 % as determined by reducing SDS-PAGE.

Background

Complement Component 1, q subcomponent (C1q) associates with C1r and C1s in order to yield the first component of

For Research Use Only

the serum complement system. Deficiency of C1q has been associated with lupus erythematosus and glomerulonephritis. C1q is composed of 18 polypeptide chains: six A-chains, six B-chains, and six C-chains. Southern blot analysis of chromosomal DNA from vertebrate species demonstrated highest similarity between the C1qB genes, followed by C1qC and finally C1qA. Sequence comparison of C1q from three different species have shown that the B chains have the strongest similarity. C1q was already present at embryonic day 14 (E14) and showed little change in abundance through six weeks postnatal. At E16, C1qB mRNA was present at high abundance in putative microglia/macrophages in cortical marginal and intermediate zones, and hippocampal analge.