A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CD200 protein (His tag)

Catalog No. PKSH031373

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms OX-2 Membrane Glycoprotein, CD200, MOX1, MOX2

Species Human

Expression Host HEK293 Cells
Sequence Met1-Gly232
Accession P41217
Calculated Molecular Weight 24 kDa
Observed molecular weight 35 kDa
Tag C-His

**Bioactivity** Testing in progress

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** Please contact us for more information.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

# **Background**

CD200 is a transmembrane immunoregulatory protein that belongs to the immunoglobulin superfamily. It contains one Ig like V type domain and one Ig like C2 type domain in its extracelluar domain. CD200 is widely but not ubiquitously

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

 $Email: \underline{tech support@elabscience.com}$ 

# **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

expressed. Its receptor (CD200R) is restricted primarily to mast cells; basophils; macrophages; and dendritic cells; which suggests myeloid cell regulation as the major function of CD200. CD200 and CD200R associate via their respective Nterminal Ig-like domains. In myeloid cells; CD200R initiates inhibitory signals following receptor-ligand contact. In T cells; CD200 functions as a co-stimulatory molecule independent of the CD28 pathway. In addition; CD200 also plays an important role in prevention of graft rejection; autoimmune diseases and spontaneous abortion.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com