

Recombinant Human Follistatin/FST Protein (Fc Tag)

Catalog No. PKSH031489

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	FS
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Trp 344
Accession	NP_037541.1
Calculated Molecular Weight	61.7 kDa
Observed molecular weight	70 kDa
Tag	C-hFc
Bioactivity	<ol style="list-style-type: none"> 1. Measured by its ability to bind human INHBA-his in a functional ELISA. 2. Measured by its ability to bind mouse INHBA-his in a functional ELISA. 3. Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED50 for this effect is typically 0.5-3 µg/mL in the presence of 10 ng/ml Recombinant Human ctivin A.

Properties

Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	<p>Lyophilized from sterile PBS, pH 7.4</p> <p>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.</p> <p>Please refer to the specific buffer information in the printed manual.</p>
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Follistatin is a single-chain gonadal protein that specifically inhibits follicle-stimulating hormone release. The single FST gene encodes two isoforms; FST317 and FST344 containing 317 and 344 amino acids respectively; resulting from alternative splicing of the precursor mRNA. In a study in which 37 candidate genes were tested for linkage and association with polycystic ovary syndrome (PCOS) or hyperandrogenemia in 150 families; evidence was found for linkage between PCOS and follistatin. Follistatin are expressed and subserve local regulatory roles in numerous extragonadal tissues; including brain; adrenal; bone marrow; and placenta but perhaps most notably in anterior pituitary- the classical target tissue for inhibin; the activin-follistatin system may play a key role in early embryogenesis. Follistatin binds directly to activin and functions as an activin antagonist. Specific inhibitor of the biosynthesis and secretion of pituitary follicle stimulating hormone follistatin is a binding protein to activin. Since activin binds to follistatin; it is imperative to determine the nature of the activin/follistatin binding complex.

For Research Use Only