## Recombinant Human sFRP1/SARP2 Protein (His Tag)

Catalog Number: PKSH031491



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Synonyms FRP;FRP-1;FRP1;FrzA;SARP2

Species Human

Expression Host HEK293 Cells
Sequence Met 1-Lys 314
Accession NP\_003003.3
Calculated Molecular Weight 34.0 kDa
Observed molecular weight 38 kDa
Tag C-His

**Bioactivity** Measured by its ability to inhibit proliferation of HeLa human cervical epithelial

carcinoma cells. The ED50 for this effect is typically 2. 5-10 µg/ml.

## **Properties**

**Purity** > 97 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 97 % as determined by reducing SDS-PAGE.

### **Background**

Secreted frizzled-related protein 1, also known as sFRP1, is a 35 kDa prototypical member of the SFRP family. SFRP family consists of five secreted glycoproteins in humansacting as extracellular signaling ligands. Each is approximately 300 amino acids in length and contains a cysteine-rich domain (CRD) that shares 30-50% sequence homology with the CRD ofFrizzled(Fz) receptors, a putative signal sequence, and a conserved hydrophilic carboxy-terminal domain.SFRPs act as soluble modulators of Wnt signaling, counteracting Wnt-induced effects at high concentrations and promoting them at lower concentrations. SFRPs are able to bindWntproteins and Fz receptors in the extracellular compartment. The

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Human sFRP1/SARP2 Protein (His Tag)

Catalog Number:PKSH031491



interaction between SFRPs and Wnt proteins prevents the latter from binding the Fz receptors. The Wnt pathway plays a key role in embryonic development, cell differentiation and cell proliferation. The deregulation of this critical developmental pathway occurs in several human tumor entities. Mouse sFRP1 is highly expressed in kidney and embryonic heart, as well as in the eye, where it is principally localized to the ciliary body and the lens epithelium.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>