

Recombinant Human FGFR1/CD331 Protein (His Tag)

Catalog No. PKSH031506

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	bFGF-R-1;BFGFR;CD331;CEK;FGFBR;FGFR-1;FLG;FLT-2;FLT2;HBGFR;HH2;HRTFDS;KAL2;N-SAM;OGD
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Glu 285
Accession	NP_075594.1
Calculated Molecular Weight	31.0 kDa
Observed molecular weight	50-55 kDa
Tag	C-His
Bioactivity	Measured by its ability to inhibit FGF acidic dependent proliferation of Balb/C 3T3 mouse fibroblasts. The ED50 for this effect is typically 0.1-0.6 µg/mL.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

FGFR1; also known as CD331; belongs to the fibroblast growth factor receptor subfamily where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. Fibroblast growth factors (FGFs) (FGF1 - 10 and 16 - 23) are mitogenic signaling molecules that have roles in angiogenesis; wound healing; cell migration; neural outgrowth and embryonic development. FGFs bind heparan sulfate glycosaminoglycans; which facilitates dimerization (activation) of FGF receptors. FGFR1 is a full-length representative protein consists of an extracellular region; composed of three immunoglobulin-like domains; a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of FGFR1 interacts with fibroblast growth factors; setting in motion a cascade of downstream signals; ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. CD331 can be detected in astrocytoma; neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines; however isoform 17; isoform 18 and isoform 19 are not detected in these cells. Defects in FGFR1 are a cause of Pfeiffer syndrome ;idiopathic hypogonadotropic hypogonadism; Kallmann syndrome type 2; osteoglophonic dysplasia and trigonocephaly non-syndromic.