

Recombinant Human TGFBI/BIGH3 Protein (His Tag)

Catalog No. PKSH031520

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	BIGH3;CDB1;CDG2;CDGG1;CSD;CSD1;CSD2;CSD3;EBMD;LCD1
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-His 683
Accession	NP_000349.1
Calculated Molecular Weight	74.0 kDa
Observed molecular weight	65 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 75 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 75 % as determined by reducing SDS-PAGE.

Background

TGFBI is an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins.

For Research Use Only

TGFBI plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. TGFBI is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in TGFBI are associated with multiple types of corneal dystrophy. TGFBI can bind to type I, II, and IV collagens. This adhesion protein may play an important role in cell-collagen interactions. In cartilage, TGFBI may be involved in endochondral bone formation. Loss of the TGFBI is sufficient to induce specific resistance.