

Recombinant Human Cathepsin C/CTSC/DPPI Protein (His Tag)

Catalog No. PKSH031574

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CPPI;DPP-I;DPP1;DPPI;HMS;JP;JPD;PALS;PDON1;PLS
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Leu 463
Accession	NP_001805.3
Calculated Molecular Weight	51.0 kDa
Observed molecular weight	55 kDa
Tag	C-His
Bioactivity	Measured by its ability to cleave the fluorogenic peptide substrate, Gly-Arg-7-amido-4-methylcoumarin (GRAMC). The specific activity is > 200 pmoles/min/μg. (Activation description: The proenzyme needs to be activated by Cathepsin L for an activated form)

Properties

Purity	> 92 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 92 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Cathepsins are proteases found in many types of cells conserved in all animals, which have a vital role in mammalian cellular turnover such as bone resorption. The lysosomal cysteine protease Cathepsin C (CTSC), also known as dipeptidyl peptidase I (DPPI/DPP1), activates a number of granule-associated serine proteases with pro-inflammatory and immune functions by removal of their inhibitory N-terminal dipeptides. This lysosomal exo-cysteine protease belonging to the peptidase C1 family. Active cathepsin C is found in lysosomes as a 200-kDa multimeric enzyme. Subunits constituting this assembly all arise from the proteolytic cleavage of a single precursor giving rise to three peptides: the propeptide, the alpha- and the beta-chains. It is a central coordinator for activation of many serine proteases in immune/inflammatory cells. Defects in the Cathepsin C have been shown to be a cause of Papillon-Lefevre disease, an autosomal recessive disorder characterized by palmoplantar keratosis and periodontitis. Cathepsin C plays a key role in the activation of several degradative enzymes linked to tissue destruction in inflammatory diseases. Thus, it is a therapeutic target for the treatment of a number of inflammatory and autoimmune diseases.

For Research Use Only