

Recombinant Human IL4RA/CD124 Protein (His Tag)

Catalog Number:PKSH031630



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-4 receptor subunit alpha;IL-4 receptor subunit alpha;IL-4R subunit alpha;IL-4R-alpha;IL-4RA;CD124;IL-4-binding protein;IL4-BP;IL4R;IL4RA
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-His 232
Accession	NP_000409.1
Calculated Molecular Weight	25.3 kDa
Observed molecular weight	43-48 kDa
Tag	C-His
Bioactivity	1. Immobilized human IL4R-His at 10 µg/mL (100 µl/well) can bind biotinylated human IL4. The EC50 of biotinylated human IL4 is 20.8-48.5 ng/mL. 2. Measured by its ability to inhibit IL-4 dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 4-20 ng/ml in the presence of 2 ng/ml IL-4.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human IL4RA/CD124 Protein (His Tag)

Catalog Number:PKSH031630



function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD124, also known as interleukin 4 receptor (IL4R), is a type I transmembrane protein that can regulate IgE antibody production in B cells through binding to interleukin 4 and interleukin 13 and promote differentiation of Th2 cells through binding to interleukin 4. The membrane-bound form of CD124 can be hydrolyzed to soluble form which can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com