

## Recombinant Human PGD protein (His tag)

Catalog No. PKSH031635

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	6-phosphogluconate dehydrogenase, Decarboxylating, PGD, PGDH, 6PGD
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Ala 483
<b>Accession</b>	P52209
<b>Calculated Molecular Weight</b>	54.6 kDa
<b>Observed molecular weight</b>	55 kDa
<b>Tag</b>	N-His & C-His
<b>Bioactivity</b>	Testing in progress

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

6-phosphogluconate dehydrogenase(PGD) is a cytoplasm-located protein; and belongs to the 6-phosphogluconate dehydrogenase family. 6PGD is the second dehydrogenase in the pentose phosphate shunt. It catalyzes the oxidative

### For Research Use Only

decarboxylation of 6-phosphogluconate to ribulose 5-phosphate and CO<sub>2</sub>; with concomitant reduction of NADP to NADPH. Mutations within the gene coding this enzyme result in 6-phosphogluconate dehydrogenase deficiency; an autosomal hereditary disease effecting the red blood cells.