

Recombinant Human PD-1/PDCD1 Protein (His Tag)

Catalog No. PKSH031642

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Programmed cell death protein 1;PDCD1;PD-1;hPD-1;CD279;SLEB2;Hsle1
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Gln 167
Accession	NP_005009.2
Calculated Molecular Weight	17.4 kDa
Tag	C-His
Bioactivity	Immobilized human PD-1 at 10 µg/ml (100 µl/well) can bind recombinant human B7-H1 / PD-L1 / Fc chimera with a linear range of 0.02-0.4 µg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Programmed cell death 1; also known as PDCD1; is a type I transmembrane glycoprotein; and is an immunoreceptor belonging to the CD28/CTLA-4 family negatively regulates antigen receptor signaling by recruiting protein tyrosine

For Research Use Only

phosphatase; SHP-2 upon interacting with either of two ligands; PD-L1 or PD-L2. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1; IL-4; IL-10 and IFN- γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition; coligation of PD1 inhibits BCR-mediated signal by dephosphorylating key signal transducer. PD1 has been suggested to be involved in lymphocyte clonal selection and peripheral tolerance; and thus contributes to the prevention of autoimmune diseases. Furthermore; PD1 is shown to be a regulator of virus-specific CD8+ T cell survival in HIV infection. As a cell surface molecule; PDCD1 regulates the adaptive immune response. Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation; cytokine production; and cytolytic function.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017