Recombinant Human IL13RA2/CD213A2 Protein (His Tag)

Catalog No. PKSH031657

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	CD213A2;CT19;IL-13R;IL13BP;IL13R
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Leu342
Accession	NP_000631.1
Calculated Molecular Weight	38 kDa
Tag	C-His
Bioactivity	Measured by its ability to inhibit IL13-dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is typically 0.02-0.08 μg/mL.

Properties

Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

Interleukin-13 receptor subunit alpha-2 (IL13RA2/IL-13RA2) is also known as also known as cluster of differentiation 213A2 (CD213A2), IL-13 receptor subunit alpha-2, IL-13R subunit alpha-2, and IL-13RA2. The IL13RA2 is often

overexpressed in brain tumors, making Il13ra2 one of the vaccine targets for immunotherapy of glioma. IL13RA2/IL-13RA2 is a cancer-associated receptor that is present in greater than 80% of High Grade Astrocytomas (HGA) and has recently been recognized as a cytokine that predisposes breast cancer cells to metastasize. Expression of IL13Rα2 was rapidly lost from the surface of transduced cells grown in culture. The loss appeared to be related to ligands present in fetal bovine serum in the medium. None of the malignant glioma cell lines cultivated in vitro and tested to date exhibited the IL13Rα2 receptor. A recombinant virus (R5111) enters cells via its interaction with the IL13Rα2 receptor in a manner that cannot be differentiated from the interaction of wild-type virus with its receptors.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com