

# Recombinant Human LEPR/CD295 Protein (His Tag)

Catalog Number:PKSH031688



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Leptin receptor;LEP-R;HuB219;OB receptor;OB-R;CD295;LEPR;DB;OBR
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Asp 839
<b>Accession</b>	NP_002294.2
<b>Calculated Molecular Weight</b>	95.0 kDa
<b>Observed molecular weight</b>	130-140 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Immobilized human Leptin at 5 µg/ml (100 µl/well) can bind human Leptin receptor with a linear range of 0.032-4.0 µg/ml.

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Leptin Receptor or CD295 belongs to the gp130 family of cytokine receptors that are known to stimulate gene transcription via activation of cytosolic STAT proteins. This protein is a receptor for leptin (an adipocyte-specific hormone that regulates body weight), and is involved in the regulation of fat metabolism, as well as in a novel hematopoietic pathway that is required for normal lymphopoiesis. Leptin Receptor/CD295 is a transmembrane catalytic receptors found on NPY/AgRP and alpha-MSH/CART neurons in hypothalamic nuclei. Leptin receptors (Ob-Rs) are coded for by one human gene that produces six different isoforms; Ob-Ra - Ob-Rf. Ob-Rs exist as constitutive dimers at

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human LEPR/CD295 Protein (His Tag)

Catalog Number:PKSH031688



physiological expression levels. Only the Ob-Rb isoform can transduce intracellular signals and does so through activation of the JAK2/STAT3, PI 3-K and MAPK signaling cascades. Activation of Ob-Rs mediates transcriptional regulation of the hypothalamic melanocortin pathway and downregulates endocannabinoid expression. Leptin acts via leptin receptors. Leptin resistance has been proposed as a pathophysiological mechanism of obesity. In obese individuals, Ob-Ra (which is involved in active transport of leptin across the blood-brain barrier) expression is downregulated and the individual may be unresponsive to leptin signals. Ob-R antagonists are of great interest in the development of pharmacological treatments for obesity. Mutations in Leptin Receptor/CD295 have been associated with obesity and pituitary dysfunction.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017