

Recombinant Human SerpinA1/A1AT Protein (His Tag)

Catalog No. PKSH031695

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Alpha-1-Antitrypsin;Alpha-1 Protease Inhibitor;Alpha-1-Antiproteinase;Serpine A1; SERPINA1;AAT;PI;A1A;A1AT;AAT;alpha1AT;MGC23330;MGC9222;PI1;PRO2 275
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Lys 418
Accession	NP_000286.3
Calculated Molecular Weight	45.7 kDa
Observed molecular weight	55-60 kDa
Tag	C-His
Bioactivity	Measured by its ability to inhibit trypsin cleavage of a fluorogenic peptide substrate, Mca-RPKPVE-Nval-WRK(Dnp)-NH ₂ (Anaspec, Catalog#27114). The IC ₅₀ value is < 3.0 nM, as measured in 100µL reaction mixture containing 1.25 ng trypsin (Sigma, Catalog#T1426), 10 µM substrate, 50 mM Tris, 10 mM CaCl ₂ , 0.15 M NaCl, pH 7.5.

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

SerpinA1, also known as Alpha-1 antitrypsin (AAT), is a prototype member of the Serpin superfamily of the serine protease inhibitors. This serine protease inhibitor blocks the protease, neutrophil elastase. Alpha-1 antitrypsin is mainly produced in the liver and acts as an antiprotease. Its principal function is to inactivate neutrophil elastase, preventing tissue damage. SerpinA1, an acute phase protein and the classical neutrophil elastase inhibitor, is localized within lipid rafts in primary human monocytes in vitro. Its association with monocytes is inhibited by cholesterol depleting/efflux-stimulating agents and oxidized low-density lipoprotein (oxLDL) and conversely, enhanced by free cholesterol.

Furthermore, SerpinA1/monocyte association per se depletes lipid raft cholesterol as characterized by the activation of extracellular signal-regulated kinase 2, formation of cytosolic lipid droplets, and a complete inhibition of oxLDL uptake by monocytes. Alpha-1 antitrypsin deficiency is a recently identified genetic disease that occurs almost as frequently as cystic fibrosis. It is caused by various mutations in the SerpinA1 gene, and has numerous clinical implications. Alpha-1 antitrypsin deficiency is an inherited disease affecting the lung and liver. In the liver, alpha-1 antitrypsin deficiency may manifest as benign neonatal hepatitis syndrome; a small percentage of adults develop liver fibrosis, with progression to cirrhosis and hepatocellular carcinoma.