

Recombinant Human PIN1/Rotamase Pin1 Protein (His Tag)

Catalog No. PKSH031711

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	DOD;UBL5
Species	Human
Expression Host	E.coli
Sequence	Met 1-Glu 163
Accession	Q13526-1
Calculated Molecular Weight	20.3 kDa
Observed molecular weight	21 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Peptidyl-prolyl cis-trans isomerase Pin1, also known as Peptidyl-prolyl cis-trans isomerase NIMA-interacting 1, Rotamase Pin1 and PIN1, peptidyl-prolyl cis/trans isomerase (PPIase), is a nucleus protein. PIN1 is a peptidyl-prolyl isomerase that

For Research Use Only

can alter the conformation of phosphoproteins and so affect protein function and/or stability. PIN1 regulates a number of proteins important for cell-cycle progression and is presumed to operate as a molecular timer of this important process. PIN1 is an essential PPIase that regulates mitosis presumably by interacting with NIMA and attenuating its mitosis-promoting activity. PIN1 displays a preference for an acidic residue N-terminal to the isomerized proline bond. Alterations in the level of PIN1 can influence hyperproliferative diseases such as cancer. PIN1 has been implicated in multiple aspects of cell cycle regulation. It has been suggested that PIN1 function is required for both normal mitotic progression and reentry into the cell cycle from quiescence. PIN1 is also a target of several oncogenic pathways and is overexpressed in human breast cancer. Its overexpression can lead to upregulation of cyclin-D1 and transformation of breast epithelial cells in collaboration with the oncogenic pathways. PIN1 plays a pivotal role in breast development and may be a promising new anticancer target. Recent data also implicate Pin1 as playing an important role in immune responses, at least in part by increasing the stability of cytokine mRNAs by influencing the protein complexes to which they bind.