

## Recombinant Human IGF1R protein (His tag)

Catalog No. PKSH031816

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	CD221 antigen, CD221, IGF1R, IGF-1R, IGF-I R, IGF-I receptor, IGFIR, IGF-IR, IGFR, insulin-like growth factor 1 receptor, JTK13
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met1-Asn932
<b>Accession</b>	P08069
<b>Calculated Molecular Weight</b>	81 & 120 & 48 kDa
<b>Observed molecular weight</b>	110 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Testing in progress

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

The insulin-like growth factor-1 receptor (IGF1R) is a transmembrane tyrosine kinase involved in several biological

### For Research Use Only

processes including cell proliferation, differentiation, DNA repair, and cell survival. This a disulfide-linked heterotetrameric transmembrane protein consisting of two  $\alpha$  and two  $\beta$  subunits, and among which, the  $\alpha$  subunit is extracellular while the  $\beta$  subunit has an extracellular domain, a transmembrane domain and a cytoplasmic tyrosine kinase domain. The IGF-I receptor is highly expressed in all cell types and tissues. Essentially all of the biological activities of IGF-I and II have been shown to be mediated via IGF-I R. IGF1R is an important signaling molecule in cancer cells and plays an essential role in the establishment and maintenance of the transformed phenotype. Inhibition of IGF1R signaling thus appears to be a promising strategy to interfere with the growth and survival of cancer cells, is now an attractive anti-cancer treatment target.