

# Recombinant Human JAML/AMICA Protein (His Tag)

Catalog Number:PKSH031859



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	AMICA;CREA7-1;CREA7-4;Gm638;JAML
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Leu 275
<b>Accession</b>	NP_001091996.1
<b>Calculated Molecular Weight</b>	30.5 kDa
<b>Observed molecular weight</b>	43 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 90 % as determined by reducing SDS-PAGE.

## Background

Junctional adhesion molecules (JAMs) are endothelial and epithelial adhesion molecules involved in the recruitment of circulating leukocytes to inflammatory sites. JAML (Junctional adhesion molecule-like), also known as AMICA1 (Adhesion molecule interacting with CXADR antigen 1), a protein related to the JAM family, is restricted to leukocytes and promotes their adhesion to endothelial cells. It contains 2 extracellular immunoglobulin-like domains, a transmembrane segment, and a cytoplasmic tail involved in activation signaling. Monocytic JAML/AMICA1 plays a critical role in regulating monocyte transendothelial migration (TEM) probably via binding to the endothelial coxsackie and adenovirus receptor (CAR) and other tight junction-associated adhesive molecules. The Expression of JAML/AMICA1 is restricted to the hematopoietic tissues with the exception of liver. JAML may function in

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human JAML/AMICA Protein (His Tag)

Catalog Number:PKSH031859



transmigration of leukocytes through epithelial and endothelial tissues. Expressed at the plasma membrane of polymorphonuclear leukocytes, JAML/AMICA1 also enhances myeloid leukemia cell adhesion to endothelial cells.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017