
Recombinant Human Glypican 5/GPC5 Protein (His Tag)

Catalog No. PKSH031890

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	GPC5;bA93M14.1;glypican proteoglycan 5;glypican-5
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Thr 554
Accession	NP_004457.1
Calculated Molecular Weight	60.5 kDa
Observed molecular weight	60.5 kDa
Tag	C-His
Bioactivity	Immobilized human GPC5 at 5 μ g/ml (100 μ l/well) can bind human bFGF with a linear ranger of 0.156-2. 5 ng/ml.
Properties	
Purity	> 92 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



> 92 % as determined by reducing SDS-PAGE.

Background

Glypican-5 (GPC5), is a cell membrane protein which belongs to the glypican family. The glypicans compose a family of

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

Elabscience®

glycosylphosphatidylinositol-anchored heparan sulfate proteoglycans that may play a role in the control of cell division and growth regulation. So far, six members (Glypican-1/GPC1, Glypican-2/GPC2, Glypican-3/GPC3, Glypican-4/GPC4, Glypican-5/GPC5, Glypican-6/GPC6) of this family are known in vertebrates. In adult, Glypican-5 is primarily expressed in the brain. It is also detected in fetal brain, lung and liver. Glypican-5 enhances the intracellular signaling of FGF2 and HGF. It alters the cellular distribution of FGF2. The properties of Glypican-5 make it an attractive target for therapeutic intervention in rhabdomyosarcomas and other tumors that amplify and/or overexpress its gene. Glypican-5 is overexpressed in lymphoma cell lines that had shown amplification. It is a likely target for amplification, and that overexpression of GPC5 may contribute to development and/or progression of lymphomas and other tumors.

For Research Use Only