

Recombinant Human CD112/Nectin-2 Protein (His Tag)(Active)

Catalog No. PKSH031940

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Poliovirus Receptor-Related Protein 2; Herpes Virus Entry Mediator B; Herpesvirus Entry Mediator B; HveB; Nectin-2; CD112; PVRL2; HVEB; PRR2
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Leu 360
Accession	NP_002847.1
Calculated Molecular Weight	36.2 kDa
Observed molecular weight	48 kDa
Tag	C-His
Bioactivity	Measured by its binding ability in a functional ELISA. Immobilized recombinant human CD112 at 20 µg/ml (100 µl/well) can bind biotinylated DNAM1 with a linear range of 0.078-2.5 µg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Storage	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
Reconstitution	Please refer to the printed manual for detailed information.

Data



Background

Cluster of Differentiation 112 (CD112), also known as poliovirus receptor related protein 2 (PVRL2 or PRR2), is a single-pass type I transmembrane glycoprotein belonging to the Immunoglobulin superfamily. CD112 protein also serves as an entry for certain mutant strains of herpes simplex virus and pseudorabies virus, and thus is involved in cell to cell

For Research Use Only

spreading of these viruses. CD112 protein has been identified as the ligand for DNAM-1 (CD226), and the interaction of CD226/CD112 protein can induce NK cell- and CD8+ T cell-mediated cytotoxicity and cytokine secretion. CD112 has been regarded as a critical component in allergic reactions, and accordingly may function as a novel target for anti-allergic therapy.