

Recombinant Rat ART4 Protein (His Tag)

Catalog No. PKSR030190

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	ART4
Species	Rat
Expression Host	HEK293 Cells
Sequence	Met1-Lys269
Accession	NP_001166980.1
Calculated Molecular Weight	28.4 kDa
Observed molecular weight	36-41 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

ADP-ribosyltransferase 4 (Dombrock blood group), also known as Mono-ADP-ribosyltransferase 4(ART4), Dombrock blood group carrier molecule and CD297, is a protein that contains a mono-ADP-ribosylation (ART) motif. It is a

For Research Use Only

member of the ADP-ribosyltransferase gene family but enzymatic activity has not been demonstrated experimentally. ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of *E. coli*. ART4 could be detected on HEL cells and erythrocytes by FACS analysis while it was absent from activated monocytes, despite the presence of ART4 mRNA in these cells. ART is also known as the carrier of the Dombrock blood group alloantigens (Do) which is glycosylphosphatidylinositol-anchored to the erythrocyte membrane.