A Reliable Research Partner in Life Science and Medicine

Recombinant Rat LILRA5 Protein (His Tag)

Catalog No. PKSR030222

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms LILRA5
Species Rat

Expression Host HEK293 Cells
Sequence Met1-Asn248
Accession D4A6Y0
Calculated Molecular Weight 27.2 kDa
Observed molecular weight 37 kDa
Tag C-His

Bioactivity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

LILRA5 is a member of the leukocyte immunoglobulin-like receptor (LIR) family. LILR are a family of receptors possessing extracellular immunoglobulin domains. They are also known as CD85, ILTs and LIR, and can exert

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

immunomodulatory effects on a wide range of immune cells. ILT-11 contains 2 Ig-like C2-type (immunoglobulin-like) domains. It can be detected n tissues of the hematopoietic system, including bone marrow, spleen, lymph node and peripheral leukocytes. Crosslink of ILT-11 on the surface of monocytes has been shown to induce calcium flux and secretion of several proinflammatory cytokines, which suggests the roles of this protein in triggering innate immune responses.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com