

Recombinant Rat CD99/MIC2 Protein (Fc Tag)

Catalog Number:PKSR030229



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD99
Species	Rat
Expression Host	HEK293 Cells
Sequence	Met1-Gly127
Accession	B4F7A5
Calculated Molecular Weight	37.6 kDa
Observed molecular weight	47 kDa
Tag	C-hFc

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD99 is a transmembrane protein expressed on most hematopoietic cells, endothelial cells and at the borders between confluent cells. CD99 is also found expressed in the development of normal ovary and testis as well as in 25 sex cord-stromal tumors, 7 epithelial neoplasms, and 6 germ cell tumors. CD99 may be a useful marker for sex cord-stromal

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Rat CD99/MIC2 Protein (Fc Tag)

Catalog Number:PKSR030229



tumors and that its degree of reactivity correlates with the degree of differentiation in Sertoli-Leydig cell tumors. Additionally, CD99 might aid in distinguishing granulosa cell tumors of the ovary from poorly differentiated carcinomas and it has been reported to be a sensitive and specific marker for Ewing's sarcoma and primitive neuroectodermal tumor.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com