# Recombinant Rat Tie2/TEK Protein (Fc Tag)

### Catalog No. PKSR030377

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Tie-2;Tie2
Species	Rat
Expression Host	HEK293 Cells
Sequence	Met 4-Leu 743
Accession	NP_001099207.1
Calculated Molecular Weight	108 kDa
Observed molecular weight	120-130 kDa
Tag	C-hFc
Bioactivity	Immobilized S1h-3C-mANGPT2 at 10 μg/mL (100 μL/well) can bind ratTEK-Fc. The EC50 of ratTEK-Fc is 0.26-0.62μg/mL.
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	

KDa	М
116	
66.2	-
45.0	_
35.0	-
25.0	-
18.4	_
14.4	-

> 95 % as determined by reducing SDS-PAGE.

## Background

TEK, or TIE-2, is an endothelial cell-specific receptor tyrosine kinase (RTK) that is known as a functioning molecule of

#### For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

# **Elabscience**®

vascular endothelial cells. TEK comprises a subfamily of RTK with TIE, and these two receptors play critical roles in vascular maturation, maintenance of integrity and remodeling. Targeted mutagenesis of both Tek and its agonistic ligand, Angiopoietin-1, result in embryonic lethality, demonstrating that the signal transduction pathways mediated by this receptor are crucial for normal embryonic development. TEK signaling is indispensable for the development of the embryonic vasculature and suggests that TEK signaling may also be required for the development of the tumor vasculature.

For Research Use Only