

Recombinant Mouse CD45/PTPRC Protein (His & GST Tag)

Catalog No. PKSM040297

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	B220;Cd45;CD45R;L-CA;loc;Ly-5;Lyt-4;T200
Species	Mouse
Expression Host	Baculovirus-Insect Cells
Sequence	Arg453-Ser1152
Accession	AAA39458.1
Calculated Molecular Weight	108.2 kDa
Observed molecular weight	110 kDa
Tag	N-His-GST
Bioactivity	<ol style="list-style-type: none"> 1. The specific activity was determined to be 12306 nmol/min/mg using p-nitrophenyl phosphate as substrate. 2. Measured by its ability to bind biotinylated Galectin-1 in a functional ELISA.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as sterile solution of 20mM Tris, 500mM NaCl, 0.5mM GSH, 3mM DTT, pH 7.4, 10% glycerol
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD

For Research Use Only

molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Protein tyrosine phosphatase, receptor type C (CD45), also known as PTPRC is a member of the protein tyrosine phosphatase (PTP) family which is known for its function to serve as signaling molecules and to regulate a variety of cellular processes such as cell proliferation, differentiation, mitotic cycle and oncogenic transformation. CD45 is found expression specifically in hemotopietic cells. CD45 consists of an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains. It serves as an essential regulator of T-cell and B-cell antigen receptor signaling through either direct interaction with components of the antigen receptor complexes or by activating various Src family kinases required for the antigen receptor signaling and it also can suppress JAK kinases.