

Recombinant Mouse Interleukin-2/IL-2 Protein

Catalog No. PKSM040308

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	IL2;Aldesleukin;Interleukin 2;Interleukin-2;IL-2;IL2;T-cell growth factor;T cell growth factor;TCGF
Species	Mouse
Expression Host	E.coli
Sequence	Ala 21-Gln 169
Accession	P04351
Calculated Molecular Weight	18.2 kDa
Observed molecular weight	18 kDa
Tag	C-His

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Interleukin-2, also known as T-cell growth factor, TCGF, Aldesleukin and IL2, is a secreted protein which belongs to the IL-2 family. Interleukin-2 / IL-2 is produced by T-cells in response to antigenic or mitogenic stimulation, this protein is

For Research Use Only

required for T-cell proliferation and other activities crucial to regulation of the immune response. Interleukin-2 / IL-2 is normally produced by the body during an immune response. Antigen binding to the T cell receptor (TCR) stimulates the secretion of Interleukin-2 / IL-2, and the expression of IL-2 receptors IL-2R. The IL-2 / IL-2R interaction then stimulates the growth, differentiation and survival of antigen-selected cytotoxic T cells via the activation of the expression of specific genes. Interleukin-2 / IL-2 can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells. The World Reference Standard for Interleukin-2 / IL-2 is produced by the National Institute of Biological Standards and Control in the UK. A recombinant form of Interleukin-2 / IL-2 for clinical use is manufactured by Chiron Corporation with the brand name Proleukin. It has been approved by the Food and Drug Administration (FDA) for the treatment of cancers (malignant melanoma, renal cell cancer), and is in clinical trials for the treatment of chronic viral infections, and as a booster (adjuvant) for vaccines. The use of Interleukin-2 / IL-2 in HIV therapy has been found to be ineffective.