

Recombinant Mouse EphA3 Protein (His Tag)

Catalog No. PKSM040313

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AW492086;Cek4;End3;ETK1;Hek;Hek4;Mek4;Tyro4
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met1-His541
Accession	NP_796047.2
Calculated Molecular Weight	60.2 kDa
Tag	C-His
Bioactivity	Immobilized mouse EPHA3-His at 10 µg/mL (100 µL/well) can bind mouse EFNA5-Fc. The EC50 of mouse EFNA5-Fc is 4. 9-11.4ng/mL.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

EPHA3 gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. The ephrin

For Research Use Only

receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. EPHA3 gene encodes a protein that binds ephrin-A ligands. EPHA3 is involved in the retinotectal mapping of neurons. It may also control the segregation but not the guidance of motor and sensory axons during neuromuscular circuit development.