

Recombinant Mouse CD200 Protein (His Tag)

Catalog No. PKSM040319

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Mox2;OX2
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Gly 232
Accession	O54901.1
Calculated Molecular Weight	24 kDa
Tag	C-His
Bioactivity	Immobilized recombinant mouse CD200 at 1 µg/ml (100ul/well) can bind mouse CD200R1/Fc with a linear range of 1. 5-200 ng/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD200 (OX-2) is a cell surface glycoprotein that imparts immune privileges by suppressing alloimmune and autoimmune responses through its receptor, CD200R, expressed primarily on myeloid cells. Signals delivered through the

For Research Use Only

CD200:CD200R axis have been shown to play an important role in the regulation of anti-tumor immunity, and overexpression of CD200 has been reported in a number of malignancies, including CLL, as well as on cancer stem cells. The role of CD200-CD200R signaling in immune regulation of the central nervous system has become a popular field of research in recent years. Many studies have shown that there is a close correlation between CD200-CD200R, microglia activation, and Parkinson's disease (PD). The ability of CD200 to suppress myeloid cell activation is critical for maintaining normal tissue homeostasis but may also enhance the survival of migratory neoplastic cells. CD200 and CD200R associate via their respective N-terminal Ig-like domains. CD200 has been characterized as an important immunoregulatory molecule, increased expression of which can lead to decreased transplant rejection, autoimmunity, and allergic disease. Elevated CD200 expression has been reported to be associated with poor prognosis in a number of human malignancies. In addition, CD200 also plays an important role in prevention of graft rejection, autoimmune diseases and spontaneous abortion.