

Recombinant Mouse IGFBP-2/IGFBP2 Protein (Fc Tag)

Catalog Number:PKSM040345



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AI255832;IBP-2;Igfbp-2;mIGFBP-2
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met1-Gln305
Accession	NP_032368.2
Calculated Molecular Weight	56.6 kDa
Tag	C-hFc

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

IGFBP-2, also known as IGFBP2, is an insulin-like growth factor-binding protein (IGFBP). IGFBPs prolong the half-life of the IGFs, control bioavailability, activity, and distribution of insulin-like growth factor (IGF) through high-affinity IGFBP/IGF complexes. Six high-affinity IGF-binding proteins (IGFBP-1 to -6) have been identified. The six IGFBPs are structurally related but encoded by distinct genes. IGFBPs have a high affinity for IGFs. Some members of the IGFBP family have been consistently shown to inhibit IGF actions by preventing them from gaining access to the IGF receptors, while others potentiate IGF actions by facilitating the ligand-receptor interaction. IGFBP-2 is overexpressed in many malignancies and is often correlated with an increasingly malignant status of the tumor, pointing to a potential involvement of IGFBP-2 in tumorigenesis. It contains 1 IGFBP N-terminal domain and 1 thyroglobulin type-1 domain. It inhibits IGF-mediated growth and developmental rates.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017