## **Recombinant Mouse VDR/NR1I1 Protein (His Tag)**

Catalog Number: PKSM040354



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Synonyms Nr1i1
Species Mouse

**Expression Host** Baculovirus-Insect Cells

Sequence Met1-Ser422

AccessionP48281Calculated Molecular Weight49.2 kDaObserved molecular weight55 kDaTagC-His

## **Properties**

**Purity** > 86 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.0, 10% glycerol


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 86 % as determined by reducing SDS-PAGE.

## **Background**

VDR (vitamin D(1,25- dihydroxyvitamin D3)receptor), also known as NR111, belongs to the NR1I family, NR1 subfamily. It is composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain. Vitamin D receptors (VDRs) are members of the NR1I family, which also includes pregnane X (PXR) and constitutive androstane (CAR) receptors, that form heterodimers with members of the retinoid X receptor family. VDRs repress expression of 1alpha-hydroxylase (the proximal activator of 1,25(OH)2D3) and induce expression of the 1,25(OH)2D3 inactivating enzyme CYP24. Also, it has recently been identified as an additional bile acid receptor alongside FXR and may function to protect gut against the toxic and carcinogenic effects of these endobiotics. VDR is expressed in the intestine, thyroid and kidney and has a vital role in calcium homeostasis. It is the nuclear hormone

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Mouse VDR/NR1I1 Protein (His Tag)

Catalog Number:PKSM040354



receptor, also called transcription factor that mediates the action of vitamin D3. Inherited mutations in the VDR gene leads to rickets.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>