Recombinant Mouse ART4/CD297 Protein (His Tag)

Catalog No. PKSM040362

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|-----------------------------|---|
| Synonyms | 4432404K01Rik;ARTC4;DO;DOK1 |
| Species | Mouse |
| Expression Host | HEK293 Cells |
| Sequence | Met1-Lys 263 |
| Accession | EDL10575.1 |
| Calculated Molecular Weight | 28.2 kDa |
| Observed molecular weight | 40-45 kDa |
| Tag | C-His |
| Bioactivity | Not validated for activity |
| Properties | |
| Purity | > 92 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per μ g of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data

| KDa | М |
|--------------|---|
| 116 | _ |
| 66.2 | - |
| 45.0 | - |
| 35.0 | _ |
| 25.0 | - |
| 18.4 14.4 | - |
| | |

> 92 % as determined by reducing SDS-PAGE.

Background

ADP-ribosyltransferase 4 (Dombrock blood group), also known as Mono-ADP-ribosyltransferase 4(ART4), Dombrock blood group carrier molecule and CD297, is a protein that contains a mono-ADP-ribosylation (ART) motif. It is a

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

member of the ADP-ribosyltransferase gene family but enzymatic activity has not been demonstrated experimentally. ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. ART4 could be detected on HEL cells and erythrocytes by FACS analysis while it was absent from activated monocytes, despite the presence of ART4 mRNA in these cells. ART is also known as the carrier of the Dombrock blood group alloantigens (Do) which is glycosylphosphatidylinosotol-anchored to the erythrocyte membrane.

For Research Use Only