Recombinant Mouse Tie2/CD202b Protein (His Tag)

Catalog Number:PKSM040370

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	AA517024;Cd202b;Hyk;STK1;Tie-2;Tie2
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met1-Lys744
Accession	Q02858
Calculated Molecular Weight	82.4 kDa
Observed molecular weight	91 kDa
Tag	C-His
Bioactivity	Measured by its binding ability in a functional ELISA. Immobilized mouse TEK-His at 10 μg/ml (100 μl/well) can bind human Ang2-Fc with a linear range of 6. 25-200 ng/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

TEK, or TIE-2, is an endothelial cell-specific receptor tyrosine kinase (RTK) that is known as a functioning molecule of vascular endothelial cells. TEK comprises a subfamily of RTK with TIE, and these two receptors play critical roles in vascular maturation, maintenance of integrity and remodeling. Targeted mutagenesis of both Tek and its agonistic ligand, Angiopoietin-1, result in embryonic lethality, demonstrating that the signal transduction pathways mediated by this receptor are crucial for normal embryonic development. TEK signaling is indispensable for the development of the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

embryonic vasculature and suggests that TEK signaling may also be required for the development of the tumor vasculature.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com