

Recombinant Mouse BTLA/CD272 Protein (Fc Tag)

Catalog No. PKSM040375

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	A630002H24
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met1-Gly176
Accession	Q7TSA3
Calculated Molecular Weight	437 kDa
Observed molecular weight	57-62 kDa
Tag	C-hFc
Bioactivity	Immobilized mouse BTLA-Fc at 10 µg/ml (100 µl/well) can bind biotinylated mouse HVEM-Fch, The EC50 of biotinylated mouse HVEM-Fch is 15.7-36.7 ng/ml.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

BTLA is a inhibitory molecule which belongs to the Ig superfamily. It down-modulates immune responses. As such, reagents that regulate the binding of BTLA to its ligand or alter BTLA signaling have significant therapeutic promise. BTLA is crucial to understand the mechanism(s) of action of these antibodies before attempting clinical applications. BTLA is not expressed by naive T cells, but it is induced during activation and remains expressed on T helper type 1 (T(H)1) but not T(H)2 cells. BTLA is a third inhibitory receptor on T lymphocytes with similarities to cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) and programmed death 1 (PD-1).