

Recombinant Mouse TSLP Protein (His Tag)

Catalog No. PKSM040390

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Thymic stromal lymphopoietin;Thymic stroma-derived lymphopoietin;Tslp;TSLP
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Glu 140
Accession	Q9JIE6
Calculated Molecular Weight	15.4 kDa
Observed molecular weight	22-27 kDa
Tag	C-His
Bioactivity	Measured by its ability to bind human IL7RA-his in functional ELISA.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Thymic stromal lymphopoietin (TSLP) is an interleukin 7 (IL-7)-like cytokine originally characterized by its ability to promote the activation of B cells and dendritic cells (DCs). Thymic stromal lymphopoietin (TSLP) is a cytokine expressed

For Research Use Only

by epithelial cells, including keratinocytes, and is important in allergic inflammation. Subsequent studies have shown that TSLP promotes T helper type 2 (TH2) cell responses associated with immunity to some helminth parasites and the pathogenesis of many inflammatory diseases, including atopic dermatitis and asthma. TSLP can promote TH2 cytokine-associated inflammation by directly promoting the effector functions of CD4+ TH2 cells, basophils and other granulocyte populations while simultaneously limiting the expression of DC-derived proinflammatory cytokines and promoting regulatory T cell responses in peripheral tissues.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017