A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse VCL/Vinculin Protein (His Tag)

Catalog No. PKSM040490

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms 9430097D22;AA571387;AI462105;AW545629

Species Mouse

Expression Host

Sequence

Met1-Gln1066

Accession

NP_033528.3

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Met1-Gln1066

NP_033528.3

118.2 kDa

118 kDa

C-His

Bioactivity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Vinculin (VCL) is a cytoskeletal protein that is closely related to both cell-matrix interactions and cell-cell junctions. VCL is a membrane-cytoskeletal protein in focal adhesion plaques that is involved in linkage of integrin adhesion molecules to

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

the actin cytoskeleton. The protein contains an acidic N-terminal domain and a basic C-terminal domain separated by a proline-rich middle segment. This protein has multi-ligand properties and has been found to interact with a number of microfilament associated proteins, such as talin, a-actinin, and paxillin, which reportedly bind to either the head or tail domains of vinculin.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com