A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse Smad5 Protein (His & GST Tag)

Catalog No. PKSM040516

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms 1110051M15Rik;AI451355;Dwf-C;Madh5;MusMLP

Species Mouse

Expression Host Baculovirus-Insect Cells

SequenceThr2-Ser465AccessionP97454Calculated Molecular Weight79.9 kDaObserved molecular weight75-88 kDaTagN-His-GST

Bioactivity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.0, 10% glycerol


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

SMAD5 is a member of the SMAD family. Members of this family mediate signal transduction by the TGF-beta/activin/BMP-2/4 cytokine superfamily from receptor Ser/Thr protein kinases at the cell surface to the nucleus.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

SMAD5 is involved in the TGF-beta signaling pathway that results in an inhibition of the proliferation of hematopoietic progenitor cells. It is also involved in cell signalling and modulates signals of bone morphogenetic proteins (BMP's). The binding of ligands causes the oligomerization and phosphorylation of the SMAD5 protein. SMAD5 is a receptor regulated SMAD (R-SMAD) and is activated by bone morphogenetic protein type 1 receptor kinase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com