

Recombinant Mouse BLMH/BLM Hydrolase Protein (His Tag)

Catalog No. PKSM040554

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AI035728;Bh;Bmh
Species	Mouse
Expression Host	E.coli
Sequence	Asn 2-Glu 455
Accession	NP_848760.1
Calculated Molecular Weight	53.3 kDa
Observed molecular weight	47 kDa
Tag	N-His
Bioactivity	Measured by its ability to hydrolyze Met-AMC. The specific activity is > 500 pmoles/min/μg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 0.15M NaCl, 10% glycerol, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The papain superfamily member bleomycin hydrolase (BLMH) is a cytoplasmic cysteine peptidase that is highly

For Research Use Only

conserved through evolution. The only known activity of the enzyme is metabolic inactivation of the glycopeptide bleomycin (BLM), an essential component of combination chemotherapy regimens for cancer. The papain superfamily member bleomycin hydrolase (BLMH) is a neutral cysteine protease with structural similarity to a 20S proteasome. Bleomycin (BLM), a clinically used glycopeptide anticancer agent. BLMH is an essential protectant against BLM-induced death and has an important role in neonatal survival and in maintaining epidermal integrity. Sequencing revealed several putative sites phosphorylated by different types of protein kinases, but no signal sequence, transmembrane domain, N-linked glycosylation site or DNA-binding motif.