# Recombinant Mouse CD200R4/CD200RLa Protein (His

# Tag)

#### Catalog Number: PKSM040571



*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Descri	

Synonyms Cd200r4;F630107N04Rik;MCD200RLa;CD200 Receptor 4

**Species** Mouse

Expression Host

Sequence

Met 1-Thr 241

Accession

NP\_997127.1

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Met 1-Thr 241

NP\_997127.1

25 kDa

50-60 kDa

C-His

## **Properties**

**Purity** > 96 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 96 % as determined by reducing SDS-PAGE.

## **Background**

Cell surface glycoprotein CD200 receptor 4, also known as Cell surface glycoprotein OX2 receptor 4, CD200 cell surface glycoprotein receptor-like 4, CD200RLa, and CD200R4, is a single-pass type I membrane protein which belongs to theCD200R family. CD200 (OX2) is a cell surface glycoprotein that interacts with a structurally related receptor (CD200R) expressed mainly on myeloid cells and is involved in regulation of macrophage and mast cell function. In mouse there are up to five genes related to CD200R with conflicting data as to whether they bind CD200. CD200R4 contains oneIg-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. CD200R4 is highly expressed in monocytes, NK cells and a subset of NKT cells. It is weakly expressed in granulocytes and B cells (at protein level). CD200R4 is also expressed in brain, lung, testis, thymus, intestine and uterus. and in bone

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Mouse CD200R4/CD200RLa Protein (His Tag)



Catalog Number:PKSM040571

marrow derived-macrophage and dendritic cells and mast cells. CD200R4 is involved in the recruitment or surface expression of the TYROBP receptor.

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>