Recombinant Mouse EpCAM/TROP-1 Protein (His Tag)

Catalog Number: PKSM040591



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Synonyms	Neurotrophic tyrosine kinase receptor-related 1;receptor tyrosine kinase-like orphan receptor 1;ROR1;tyrosine-protein kinase transmembrane receptor ROR1;Epithelial cell adhesion molecule;Tumor-associated calcium signal transducer 1;TROP1;CD32 6;EGP;EGP-2;Egp314;Ep-CAM;EpCAM1;GA733-2;gp40;Ly74;Tacsd1;Tacstd1
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Thr 266
Accession	NP_032558.2
Calculated Molecular Weight	29 kDa
Observed molecular weight	37 kDa
Tag	C-His
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	

> 95 % as determined by reducing SDS-PAGE.

45.0 35.0 25.0 18.4 14.4

Background

Epithelial Cell Adhesion Molecule (EpCAM), also known as GA733-2 antigen, is a type â...transmembrane glycoprotein composed of an extracellular domain with two EGF-Like repeats and a cystenin-rich region, a transmembrane domain and a cytoplasmic domain. It modulates cell adhesion and proliferation. Its overexpression has been detected in many epithelial tumours and has been associated with high stage, high grade and a worse survival in some tumour types. EpCAM has been shown to function as a calcium-independent homophilic cell adhesion molecule that does not exhibit

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number: PKSM040591



any obvious relationship to the four known cell adhesion molecule superfamilies. However, recent insights have revealed that EpCAM participates in not only cell adhesion, but also in proliferation, migration and differentiation of cells. In addition, recent study revealed that EpCAM is the Wnt-beta-catenin signaling target gene and may be used to facilitate prognosis. It has oncogenic potential and is activated by release of its intracellular domain, which can signal into the cell nucleus by engagement of elements of the wnt pathway.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com