Recombinant Mouse CD34 Protein (His Tag)

Catalog Number: PKSM040592



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Trote: Centinuge before opening to ensure | |
|---|--|
| Description | |
| Synonyms | AU040960 |
| Species | Mouse |
| Expression Host | HEK293 Cells |
| Sequence | Met 1-Thr 287 |
| Accession | NP_001104529.1 |
| Calculated Molecular Weight | 28.6 kDa |
| Observed molecular weight | 50-60 kDa |
| Tag | C-His |
| Bioactivity | Measured by the ability of the immobilized protein to support the adhesion of the HUVEC human umbilical vein endothelial cell line. When 4 x 10^4 cells/well are added to mouse CD34 coated plates (0.8 µg/ml, 100 µl/well), approximately > 30 % will adhere after one hour at 37 °C. |
| Properties | |
| Purity | > 98 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per μ g of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |
| Data | |

| KDa | MK | | R | |
|------|----|---|---|---|
| 116 | | - | | |
| 66.2 | - | | | |
| 45.0 | - | 1 | | 1 |
| 35.0 | - | • | | |
| 25.0 | - | • | | |
| 18.4 | - | | | |
| 14.4 | - | | | |
| | | | | |

> 98 % as determined by reducing SDS-PAGE.

Background

Cluster of Differentiation 34 (CD34) is a member of a family of single-pass transmembrane sialomucin proteins, and may function as a cell-cell adhesion factor. CD34 protein is selectively expressed on hematopoietic progenitor cells and the small vessel endothelium of a variety of tissues. It has been widely used as a stem and progenitor cell marker, and clinical CD34+ stem cell transplantation (CD34+SCT) has been performed for tumor purging. CD34 monoclonal antibodies are

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number:PKSM040592



widely used to identify and isolate hemopoietic progenitors and to classify acute and chronic leukemias.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: www.elabscience.com Email: techsupport@elabscience.com