

Recombinant Mouse Peroxiredoxin 1/PRDX1 Protein (His Tag)

Catalog No. PKSM040618

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	MSP23;NkefA;OSF-3;OSF3;PAG;Paga;PrdxI;prx1;PrxI;Tdpx2;TDX2;TPxA
Species	Mouse
Expression Host	E.coli
Sequence	Met 1-Lys 199
Accession	P35700
Calculated Molecular Weight	23.5 kDa
Observed molecular weight	27 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, 10% glycerol, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

Peroxiredoxin-1, also known as Thioredoxin peroxidase 2, Natural killer cell-enhancing factor A, PRDX1, and PAGA, is a member of the ahpC/TSA family. Peroxiredoxin-1 is constitutively expressed in most human cells. It is induced to

For Research Use Only

higher levels upon serum stimulation in untransformed and transformed cells. Peroxiredoxins (PRDXs) are a family of antioxidant enzymes that are also known as scavengers of peroxide in mammalian cells. The overexpression of Peroxiredoxin-1, which is one of the peroxiredoxins that is a ubiquitously expressed protein, was related to a poor prognosis in several types of human cancers. Peroxiredoxin-1 is involved in redox regulation of the cell. It reduces peroxides with reducing equivalents provided through the thioredoxin system but not from glutaredoxin and may play an important role in eliminating peroxides generated during metabolism. Peroxiredoxin-1 Might participate in the signaling cascades of growth factors and tumor necrosis factor- α by regulating the intracellular concentrations of H_2O_2 . The reduced Peroxiredoxin-1 expression is an important factor in esophageal squamous cancer progression and could serve as a useful prognostic marker.