

Recombinant Mouse Peroxiredoxin 5/PRDX5 Protein (His Tag)

Catalog No. PKSM040619

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AOEB166;AOPP;Pmp20;Prdx6;PrxV
Species	Mouse
Expression Host	E.coli
Sequence	Met 49-Leu 210
Accession	P99029-1
Calculated Molecular Weight	18.5 kDa
Observed molecular weight	18.5 kDa
Tag	N-His
Bioactivity	Measured by its ability to reduce H ₂ O ₂ . The specific activity is > 100 pmoles/min/μg.

Properties

Purity	> 92 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.5 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 92 % as determined by reducing SDS-PAGE.

Background

Peroxiredoxin-5, also known as Alu corepressor 1, Antioxidant enzyme B166, Liver tissue 2D-page spot 71B,

For Research Use Only

Peroxisomal antioxidant enzyme, Thioredoxin peroxidase PMP20, Thioredoxin reductase, PRDX5 and ACR1, is cytoplasm protein which belongs to the peroxiredoxin 2 family. Peroxiredoxin-5 / PRDX5 reduces hydrogen peroxide and alkyl hydroperoxides with reducing equivalents provided through the thioredoxin system. Peroxiredoxin-5 / PRDX5 is involved in intracellular redox signaling. The Peroxiredoxins / Prx are a family of 25 kDa peroxidases that can reduce H₂O₂ using an electron from thioredoxin (Trx) or other substances. The mammalian Peroxiredoxins / Prx family is divided into six groups (PRDX1, PRDX2, PRDX3, PRDX4, PRDX5, PRDX6) on the basis of homology of amino acid sequences. They are located in the cytosol and play a role in the cell signaling system. All six mammalian peroxiredoxins are expressed in the lung. Peroxiredoxins / Prx is overexpressed in breast cancer tissues to a great extent suggesting that Peroxiredoxins / Prx has a proliferative effect and may be related to cancer development or progression.