

# Recombinant Mouse Carboxylesterase-2/CES2 Protein (His Tag)

Catalog No. PKSM040639

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms ces2A3
Species Mouse

Expression Host

Sequence

Met 1-Lys 557

Accession

NP\_663578.1

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Met 1-Lys 557

NP\_663578.1

Calculated Molecular Weight

60.4 kDa

52 kDa

C-His

**Bioactivity** Measured by its ability to hydrolyze pnitrophenylacetate. The specific activity is >

90, 000 pmoles/min/µg.

# **Properties**

**Purity** > 85 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per μg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

# Data



>85~% as determined by reducing SDS-PAGE.

# **Background**


Carboxylesterase 2 (CES2) is a member of the carboxylesterase family and belongs to the multigene family.

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Carboxylesterase 2 is responsible for the hydrolysis of ester- and amide-bond-containing drugs such as cocaine and beroin. It also serves to hydrolyze long-chain fatty acid esters and thioesters. It is speculated that carboxylesterases may play a role in lipid metabolism and the blood-brain barrier system and together with isform 1, are a serine esterase involved in both drug metabolism and activation. Human carboxylesterase 2 is commonly expressed in tumor tissues and irinotecan, a topoisomerase I inhibitor commonly used in the treatment of many solid tumors.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com