# **Recombinant Mouse GPNMB/Osteoactivin Protein (His**

# Tag)

#### Catalog Number: PKSM040661



Note: Centrifuge before opening to ensure complete recovery of vial contents.

<b>T</b>	4.0
Descri	ntion
DCSCII	DOTOIL

Synonyms DC-HIL;Dchil;ipd

**Species** Mouse

Expression Host HEK293 Cells
Sequence Met1-Asn502
Accession Q99P91
Calculated Molecular Weight 54.9 kDa
Observed molecular weight 96 kDa

Tag C-His

### **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 90 % as determined by reducing SDS-PAGE.

## **Background**

GPNMB belongs to the PMEL / NMB family, also known as Osteoactivin and Hematopoietic growth factor-inducible neurokinin 1 ( HGFIN ), is a transmembrane glycoprotein that is expressed in numerous cells, including osteoclasts, macrophages, dendritic cells, and tumor cells. It is suggested to influence osteoblast maturation, cell adhesion and migration. GPNMB protein acts as a downstream mediator of BMP-2 effects on osteoblast differentiation and function. GPNMB participates in bone mineralization, and functions as a negative regulator of inflammation in macrophages. Osteoactivin is expressed at high levels in normal and inflammatory liver macrophages suggesting a significant role in acute liver injury. The early-phase upregulation of Osteoactivin expression in the tubular epithelium in response to renal injury might play a role in triggering renal interstitial fibrosis via activation of matrix metalloproteinase expression and

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# **Recombinant Mouse GPNMB/Osteoactivin Protein (His Tag)**



Catalog Number:PKSM040661

collagen remodeling in rats. Osteoactivin as a protein that is expressed in aggressive human breast cancers and is capable of promoting breast cancer metastasis to bone.

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>