

# Recombinant Mouse JAM3/JAM-C Protein (His Tag)

Catalog Number:PKSM040667



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	1110002N23Rik;JAM-3;JAM-C;Jcam3
<b>Species</b>	Mouse
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Asn 241
<b>Accession</b>	NP_075766.1
<b>Calculated Molecular Weight</b>	25 kDa
<b>Observed molecular weight</b>	30-35 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 94 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 50mM Tris-Citrate, 300mM NaCl, pH 6.5 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 94 % as determined by reducing SDS-PAGE.

## Background

Junctional Adhesion Molecule C Protein & Antibody (JAM-C, JAM3 Protein) also known as Junctional adhesion molecule 3, JAM3, is a single-pass type I membrane protein that belongs to the immunoglobulin superfamily. It is an adhesion molecule expressed by endothelial cells (ECs) that plays a role in tight junction formation, leukocyte adhesion, and transendothelial migration. JAM-C is an adhesion molecule that is expressed on cells within the vascular compartment and epithelial cells and, to date, has been largely studied in the context of inflammatory events. JAM-C is also expressed in peripheral nerves and that this expression is localized to Schwann cells at junctions between adjoining myelin end loops. JAM-C was recently shown to be a counter receptor for the leukocyte beta2-integrin Mac-1 (CD11b/CD18), thereby mediating interactions between vascular cells, particularly in inflammatory cell recruitment. JAM-C is up-

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Mouse JAM3/JAM-C Protein (His Tag)

Catalog Number:PKSM040667



regulated by oxidized low-density lipoprotein (LDL) and may thereby contribute to increased inflammatory cell recruitment during atherosclerosis. JAM-C may therefore provide a novel molecular target for antagonizing interactions between vascular cells in atherosclerosis. JAM-C was shown to undergo a heterophilic interaction with the leukocyte beta2 integrin Mac-1, thereby mediating interactions between vascular cells in inflammatory cell recruitment. The homophilic interaction of JAM-C can mediate tumor cell-endothelial cell interactions and may thereby be involved in the process of tumor cell metastasis.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)