

Recombinant Mouse CD147/Basigin Protein (His & Fc Tag)

Catalog No. PKSM040728

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Basigin;Basic immunoglobulin superfamily;HT7 antigen;Membrane glycoprotein gp42;CD147;Bsg;AI115436;AI325119;EMMPRIN;HT-7
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Arg 209
Accession	NP_001070652.1
Calculated Molecular Weight	48.6 kDa
Observed molecular weight	60-70 kDa
Tag	C-His-Fc
Bioactivity	Using the Octet RED System, the affinity constant (Kd) of mouse CD147-Fc bound to Human PPIA-His was 0.4 μ M.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

CD147/EMMPRIN (Extracellular Matrix Metalloproteinase Inducer), also known as Basigin (BSG), is a transmembrane glycoprotein with different forms resulted from different modes of glycosylation and N-terminal sequence variants. It is a member of the immunoglobulin superfamily with homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. This protein play important roles in variety of events including spermatogenesis, embryo implantation, neural network formation. CD147 induces the production and release of matrix metalloproteinases (MMP) in the surrounding mesenchymal cells and tumor cells, and thereby promotes invasion, metastasis, growth and survival of malignant cells. Furthermore, CD147 also serves as a receptor for extracellular cyclophilinthe and its association with integrins might be important in signal transduction. Recently, CD147 displays increased expression in many cancers, and it has been previously demonstrated to participate in cancer metastasis and progression. Thus, CD147 and its antibody are used as an effective treatment for malignant cancers.