Recombinant Mouse S100A1 Protein (His Tag)(Active)

Catalog No. PKSM040770

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	AI266795, S100, S100a
Species	Mouse
Expression Host	E.coli
Sequence	Gly 2-Ser 94
Accession	P56565
Calculated Molecular Weight	12.5 kDa
Observed molecular weight	14 kDa
Tag	N-His
Bioactivity	Measured by its ability to bind human AGER in a functional ELISA.
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data

KDa	М
116	
66.2	-
45.0	-
35.0	
25.0	-
18.4	-
14.4	

> 95 % as determined by reducing SDS-PAGE.

Background

S100A1 is a Ca2+binding protein of the EF-hand type that belongs to the S100 protein family. S100 proteins consisting of at least 19 members exist as dimers in the cytoplasm and/or nucleus of a wide range of cells, and are involved in the

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

regulation of a number of cellular processes such as cell-cycle progression and cell differentiation. This protein has been shown to function in the processes including stimulation of Ca2+-induced Ca2+ release, inhibition of microtubule assembly, and inhibition of PKC-mediated phosphorylation.. Phosphoglucomutase is a target protein whose activity is antagonistically regulated by S100A1, and recently, S100A1 is also identified as a potent molecular chaperone and a new member of the Hsp70/Hsp90 multichaperone complex. S100A1 displays a tissue-specific expression pattern with highest levels in myocardium and is considered to be an important regulator of cardiac contractility. Accordingly, reduced expression or mutations of S100A1 gene have been implicated in cardiomyopathies.

For Research Use Only